

~~N66 24691~~

X-611-66-57

NASA TM X-55458

ON THE PRODUCTION OF LOW-ENERGY COSMIC-RAY ELECTRONS

GPO PRICE \$ _____

CFSTI PRICE(S) \$ _____

BY

P. B. ABRAHAM

Hard copy (HC) 3.00

K. A. BRUNSTEIN

Microfiche (MF) .75

T. L. CLINE

853 July 65

FACILITY FORM 602

N66 24691
(ACCESSION NUMBER)

(THRU)

75
(PAGES)

(CODE)

TMX-55458
(NASA CR OR TMX OR AD NUMBER)

29
(CATEGORY)

FEBRUARY 1966



GODDARD SPACE FLIGHT CENTER
GREENBELT, MARYLAND

Energetic Particles Preprint Series

ON THE PRODUCTION OF
LOW-ENERGY COSMIC-RAY ELECTRONS

by
P. B. Abraham
K. A. Brunstein
T. L. Cline

February 1966

GODDARD SPACE FLIGHT CENTER
Greenbelt, Maryland

ON THE PRODUCTION OF
LOW-ENERGY COSMIC-RAY ELECTRONS

by
P. B. Abraham, K. A. Brunstein*, and T. L. Cline
Goddard Space Flight Center, Greenbelt, Maryland

Abstract

The production of cosmic-ray electrons of characteristically low energies is investigated. Secondary sources, other than that of meson decay, are considered, and constraints are placed on both secondary and primary sources.

(1) (Calculations are made of the intensity of low-energy knock-on and beta-decay electrons which are secondary to cosmic-ray interactions.) In particular, knock-on production is calculated in the 100 KeV to 50 BeV kinetic-energy interval. Interstellar losses due to ionization, leakage from the galaxy, synchrotron, bremsstrahlung, and inverse compton effects are considered, as well as those due to plasma excitation, the red-shift, bremsstrahlung, synchrotron, and inverse compton effects in the intergalactic medium.) It is shown that the intensity of low-energy relativistic electrons from these sources is not negligible compared with that from the $\pi\text{-}\mu\text{-}e$ process, but does not account for the observed interplanetary electron intensity.

(2) (Energy inputs to the injected secondary electrons by a possible solar electric field of low magnitude, and by a possible galactic Fermi acceleration

*NAS-NASA Post-Doctoral Resident Research Associate.

are investigated. It is shown that one such input is necessary if the observed low-energy interplanetary electron intensity is to be attributed to secondary production.) A heliocentric field which does allow for a fit to the low-energy data cannot, however, account for the high-energy BeV electrons found to be in excess of those from $\pi^- \mu^- e$ production. The Fermi acceleration shown to be necessary to provide a fit is greater than that usually postulated for cosmic-ray protons, and also requires that the ratio of escape losses to acceleration, λ/α , be much smaller than is usually assumed for protons. This distinction is acceptable only if one postulates a significant difference between interstellar proton and electron propagation.

(3) The observation that the velocity spectrum of electrons in the energy per-unit-mass region of 7 to 25 closely approximates that of the cosmic-ray protons, and the necessity of constraints on the secondary electron hypothesis outlined above, suggest that most of the low-energy electrons are of primary origin. The similarity between this conclusion and the conclusion (based on the measurement of the charge ratio of electrons) that the higher-energy electrons are mostly primary is discussed.

TABLE OF CONTENTS

	<u>Page</u>
Abstract	iii
I. INTRODUCTION AND DISCUSSION OF OBSERVATIONS	1
II. SECONDARY GALACTIC ELECTRONS	6
a. <u>Knock-on Electrons</u>	6
b. <u>Neutron-Decay Electrons</u>	11
c. <u>Galactic Energy-Loss Mechanisms</u>	15
d. <u>Calculation of Secondary Galactic Electron Intensity</u>	19
III. SECONDARY METAGALACTIC ELECTRONS	23
a. <u>Production Mechanisms</u>	23
b. <u>Metagalactic Energy-Loss Mechanisms</u>	24
c. <u>Calculation of Secondary Metagalactic Electron Intensity</u>	27
IV. ENERGY INPUTS TO SECONDARY ELECTRONS	27
a. <u>Heliocentric Electric Field</u>	28
b. <u>Galactic Fermi Acceleration</u>	30
V. CONCLUSION	34
APPENDIX	39
a. <u>Predominance of Energy Increase</u>	40
b. <u>Predominance of Energy Decrease</u>	46
REFERENCES	50
FIGURES	53

ON THE PRODUCTION OF LOW-ENERGY COSMIC-RAY ELECTRONS

I. INTRODUCTION AND DISCUSSION OF OBSERVATIONS

The study of cosmic radiation has been, for the most part, the measurement of the intensities and energy spectra of the protons and other nuclei which possess the bulk of the cosmic-ray energy content. Recently, the electromagnetic component began to be investigated: Earl (1961) and Meyer and Vogt (1961) found electrons which have proton-like energies, but which, in the BeV region, have only a small fraction of the proton intensity. Also, DeShong, Hildebrand, and Meyer (1964) later found that the electron flux is partially composed of positrons. In addition, Kraushaar et al. (1965) set a new upper limit to the high-energy gamma-ray intensity. Since one can assume that some of these electrons and gamma rays may be primary cosmic rays and some may be secondary to cosmic-ray interactions, it follows that a great deal of new information about the origins and behavior of cosmic rays in the galaxy might come from a study of these rarer components. Most recently, a component of relativistic electrons in the few-MeV energy region was found in interplanetary space by Cline, Ludwig, and McDonald (1964). If we assume, as a working hypothesis, that the electrons of these extremely low energies are also of cosmic origin, we may learn something more by comparing their properties with those of the higher-energy electrons, protons and other cosmic rays. It is the purpose of this paper to discuss the possible sources of these particles.

Cosmic rays are presumably created when very low-energy particles are injected into some region where they are then accelerated; if this region is not interstellar space itself, they may temporarily be stored in or near that region before propagating through the galaxy; finally they are modulated in the solar environment before being detected. Certainly, several processes may compete for production, and others may take place in uncertain chronological order. The most fundamental model of interstellar acceleration, introduced by Fermi (1949), and the most quantitative model of solar modulation, described by Parker (1963), and combinations and variations of these, may possibly describe a great share of cosmic-ray origin and propagation. Since Fermi's model depends only on the particle's total energy per unit mass, which is a function of velocity (independent of charge or mass), and since Parker's model also depends on velocity alone at asymptotically low particle rigidity, we choose to represent all quantities in terms of the total energy per unit mass, γ .

The first measurements of the low-energy electrons are shown in Figure 1, in which the intensity, $(dJ/d\gamma)_e$, is plotted versus γ . These particles were found in interplanetary space with Explorer XVIII. The argument supporting a non-local origin of the electrons, put forth by the observers, rests on the fact that they undergo systematic intensity modulations of greater magnitude than any possible cosmic-ray parent component. The actual intensity is uncertain, since some fraction of the steady or baseline intensity may be local or instrumental; thus, the true intensity is between the magnitude of a typical modulated increase and the total magnitude of baseline plus the increase. Both the steady and incremental intensities are shown in the Figure; a typical increase is apparently energy-independent and is about 50 percent the steady value.

Also shown are the electron measurements in the higher-energy region; not all observations known are included, but those are displayed that are the only data available in a given energy interval (Schmoker and Earl, 1965, Agrinier et al., 1964, Daniel and Stephens, 1965), the most recent data with the best statistics (L'Heureux and Meyer, 1965), and the original data (Earl, 1961, Meyer and Vogt, 1961). It is inferred from the measurements on the positron to electron ratio (DeShong, Hildebrand and Meyer, 1964, Hartman, Meyer and Hildebrand, 1965) that approximately one third to two thirds of the electrons in the 1 BeV region are from a source other than π - μ -e decay. Since we can expect that the π - μ -e differential spectrum peaks in the $20 < \gamma < 200$ region, all of the very low-energy electrons must be from another source. The more recently obtained higher-energy data can be fitted to a common curve, but it departs from the low-energy fit, shown in Figure 1 as a dashed line, by an amount between one and two orders of magnitude at the high energies, where $\gamma > 1000$. Whether, after the π - μ -e component has been subtracted out, the remaining spectrum is of one smooth form, or is the sum of two or more separate spectra, cannot be determined until after the differential intensities in the $10 < \gamma < 100$ and $\gamma > 10^4$ regions are known and after the π - μ -e component itself is accurately determined by calculation and measurement of the e^+/e^- ratio. The sources of the electrons of $\gamma \lesssim 50$ most likely pose a quite separate question from that of the higher-energy electrons. The very low-energy electrons are shown again in Figure 2a compared with a representation of the composite primary cosmic-ray proton intensity, observed by McDonald and Ludwig (1964), and the primary alpha-particle intensity, observed by Fan, Gloeckler, and Simpson (1965). These electron, proton and alpha measurements, made on the same satellite,

are shown together with balloon measurements by Balasubrahmanyam and McDonald (1964) and by Fichtel, Guss, Kniffen, and Neelakantan (1964). The proton and alpha intensities fit a common curve when the alpha values are multiplied by a constant factor of ≈ 5 , to take into account the relative abundances. This composite representation of the 1963 proton and alpha spectra, which was introduced recently by Balasubrahmanyam, Boldt, and Palmeira (1965) fits the data all the way through this low-energy region; adapted to our units, it is

$$\frac{dJ}{d\gamma} = .35 (\gamma - 1)^{1.5} (\gamma - .47)^{-4} \text{ particles/cm}^2 \text{ sec-sr-unit } \gamma.$$

The similarity between the observed electron and proton differential velocity spectra has been discussed in a paper by Brunstein and Cline (1965), from which this figure was taken. They point out that after solar demodulation is taken into account by use of the model of Parker (1963) the electrons and protons fit even more closely the same spectrum, as shown in Figure 2b. The fact that this spectrum is a simple power law in total energy, which, for the protons at least, is continuous from a nonrelativistic γ of 1.02 up to the extreme relativistic region, is suggestive of a Fermi acceleration. However, the fact that this fit is accomplished before the usual correction for the 2.5 g cm^{-2} galactic path length is made, means that either it is accidental or that electrons and protons are indeed propagated in a spectrally neutral fashion. As outlined by Brunstein and Cline (1965), several possibilities present themselves; low-energy electrons and protons may be Fermi-accelerated from neutral material in the galactic medium in such a manner as to over-compensate ionization losses, or the 2.5 g cm^{-2}

path with its attendant spectral alteration and fragmentation probabilities may apply only to the heavies, or the electrons and protons may have a metagalactic origin, in which case during their travel to the solar system they encounter very little material. It is clear that these questions cannot at present be answered. What we will investigate in this paper is the possibility that the low-energy electrons are secondary to galactic cosmic rays, and the relation of this possibility to the question of a primary electron source.

Hayakawa and Okuda (1962) originally suggested that high-energy electrons should arise from $\pi \rightarrow \mu \rightarrow e$ decay following meson production in collisions of cosmic rays with the interstellar gas. Their calculations, and those of Jones (1963), Pollack and Fazio (1963), Ginzburg and Syrovatskii (1964), Gould and Burbidge (1965), and others showed that indeed a certain fraction of the observed electrons of $\gamma \gtrsim 50$ may originate in this manner. The high-energy positron to electron ratio, found by DeShong, et al. (1964) and by Hartman, et al. (1965) to be low, indicates that less than half of the observed electrons in the $\gamma \gtrsim 200$ region are secondaries from this process. Brunstein (1965) showed that Coulomb collisions could similarly account for only a fraction of the observed electrons with $5 < \gamma < 50$ unless a Fermi-like post-acceleration was incorporated. The majority of electrons in all observed energy regions are therefore of tentatively unexplained origin. In the calculations that follow we investigate in detail some of the various production mechanisms which may be responsible.

II. SECONDARY GALACTIC ELECTRONS

a. Knock-on Electrons

The most certain process by which cosmic rays can produce low-energy relativistic electrons through interactions with the interstellar medium is that of Coulomb collisions. In this process, energy is transferred to an atomic electron in great excess of its binding energy, and it recoils in billiard-ball fashion; free plasma electrons are turned into cosmic-ray secondaries in the same way. These knock-on electrons are thus a necessary consequence of the traversal of cosmic rays through the galactic medium.

We use the cross section for knock-on production calculated by Bhabha (1938). The particular form of the cross section incorporated is that for spin $\frac{1}{2}$, in order to obtain quantitatively accurate results to very high energies, $\gamma \approx 10^5$ or 10^6 . The form used by Brunstein (1965) was for spin 0 but was sufficiently accurate for the $5 \lesssim \gamma \lesssim 50$ region. The differential cross section for the production of an electron having total energy per unit mass in $d\gamma_e$ at γ_e by the collision of a cosmic ray of particle species j having normalized energy γ_j with a target of charge Z_I and atomic number A_i is

$$\Phi_i(\gamma_e, \gamma_j) d\gamma_e = \left[\frac{2\pi N_o Z_i r_e^2 Z_j^2}{A_i (1 - \gamma_j^{-2})} \left\{ \frac{1}{(\gamma_e - 1)^2} - \frac{s \left(\gamma_j + \frac{s + 1/s}{2} \right)}{(\gamma_e - 1) \gamma_j^2} + \frac{s^2}{2\gamma_j^2} \right\} \right] d\gamma_e$$

cm²/gram, (1)

in which N_o is Avogadro's number, r_e is the classical radius of the electron:

$r_e = e^2/mc^2 = 2.82 \times 10^{-13}$ cm, and $s = m_e/m_p \approx 1/1836$, and for which the maximum transferable energy is

$$\gamma_{\max} = 1 + (\gamma_j^2 - 1) / \left(s \left\{ \gamma_j + \frac{s + 1/s}{2} \right\} \right) \quad (2)$$

We make the usual approximation that the elemental abundances in the interstellar medium relative to hydrogen, a_i , can be represented by taking that of helium to be about 0.10 and by ignoring the higher-Z components. This approximation introduces a negligible error, particularly since the cross section varies only as Z_i/A_i to the first power. The contributions by the various nuclear species in the cosmic radiation are more important, varying as the square of Z_j . Using the cosmic-ray abundances relative to protons, b_j , given by Ginzburg and Syrovatskii (1961), we estimate that the total contribution from primaries of charge $Z_j \geq 2$ will be about an additional 0.75. Thus

$$\Phi(\gamma_j) = \sum_i \frac{a_i Z_i}{A_i} \sum_j b_j Z_j^2 \Phi_i(\gamma_j) \approx 1.75 \Phi_1(\gamma_p)$$

The source density of electrons due to galactic knock-on production is given by the following integral,

$$Q(\gamma_e, \vec{r}) = 1.75 \rho(\vec{r}) 4\pi \int_{\gamma_1}^{\gamma_2} \Phi_1(\gamma_e, \gamma_p) \frac{dJ(\vec{r})}{d\gamma_p} d\gamma_p \quad (3)$$

electrons/cm³-sec-unit γ ,

in which $\rho(\vec{r})$ is the interstellar density in g/cm^3 as a function of galactic position \vec{r} , $\frac{dJ(\vec{r})}{d\gamma_p}$ is the differential cosmic-ray proton intensity in particles/ $\text{cm}^2\text{-sec-sr-unit } \gamma$, and in which the limits of integration are functions of electron energy $\gamma_{1,2} = \gamma_{1,2}(\gamma_e)$.

We ignore any possible dependence on galactic position, \vec{r} , and use the cosmic-ray proton spectrum discussed by Brunstein and Cline (1966), which is a smooth fit to the high-energy spectrum, summarized by Ginzburg and Syrovatskii (1964), to the intermediate energy spectrum (McDonald and Webber, 1961) and to the solar demodulated low-energy proton spectrum:

$$dJ/d\gamma_p = 1.1 \gamma_p^{-2.5} \text{ protons}/\text{cm}^2\text{-sec-sr-unit } \gamma. \quad (4)$$

In order to evaluate the limits of integration in equation (3), we use the constraint on the cross section expressed by equation (2). Since the maximum possible energy is unlimited, $\gamma_2 = \infty$, while γ_1 is determined by solving the inequality $\gamma_e \leq \gamma_{\text{max}}$ for γ_p . The result is that $\gamma_p \geq \gamma_1$, where

$$\gamma_1 = \frac{s}{2}(\gamma_e - 1) + \left\{ 1 + \frac{1}{2}(1 + s^2)(\gamma_e - 1) + \frac{s^2}{4}(\gamma_e - 1)^2 \right\}^{1/2}.$$

It can be shown that γ_1 is greater than the largest root of $\Phi_1(\gamma_e, \gamma_p)$ as a function of γ_p , which means that Φ_1 is positive throughout the range of integration. Performing the integral of equation (3), we obtain

$$\begin{aligned}
Q(\gamma_e) = & 1.75(4\pi\rho)(1.1)\left(\frac{2\pi N_o r_e^2}{(\gamma_e - 1)^2}\right) \cdot \left\{ \frac{2}{3} \gamma_1^{-\frac{3}{2}} \right. \\
& + \left[1 + \frac{s^2(\gamma_e - 1)^2}{2} - \frac{(1 + s^2)(\gamma_e - 1)}{2} \right] \cdot \left[-\frac{\pi}{2} - \frac{2}{3} \gamma_1^{-\frac{3}{2}} - \frac{1}{2} \ln \left| \frac{\gamma_1^{\frac{1}{2}} - 1}{\gamma_1^{\frac{1}{2}} + 1} \right| + \arctan \gamma_1^{\frac{1}{2}} \right] \\
& \left. - s(\gamma_e - 1) \cdot \left[\frac{\pi}{2} - 2\gamma_1^{-\frac{1}{2}} - \arctan(\gamma_1^{\frac{1}{2}}) - \frac{1}{2} \ln \left| \frac{\gamma_1^{\frac{1}{2}} - 1}{\gamma_1^{\frac{1}{2}} + 1} \right| \right] \right\} \quad (5)
\end{aligned}$$

electrons/cm³-sec-unit γ ,

in which s and γ_1 are given above.

Since the cross section is valid over the range $1 < \gamma_e < 10^6$ and the cosmic-ray spectrum (which strongly depends on the modulation model only in the region $\gamma_p < 2$ where the electron production is of minor importance) is reliable over the region $\gamma_p < 10^6$, the resulting source strength $Q(\gamma_e)$ is valid over that entire region of γ_e . An approximation to the expression of equation (5), which allows for mathematical simplifications and gives an excellent numerical fit in the $1.2 < \gamma_e < 10^6$ region of physical interest is plotted in Figure 3. It is

$$Q(\gamma_e) = 4.9 \rho (\gamma_e - 1)^{-2.76} \text{ electrons/cm}^3\text{-sec-unit } \gamma. \quad (6)$$

In order to check this result we can instead integrate the cross section over the proton spectrum of McDonald and Webber (1961). The form they fit to the observations is $J(>R) = 0.40 R^{-1.25}$ protons/cm²-sec-sr, in which the rigidity R is measured in BV. We differentiate this spectrum and change the units to obtain

$$\frac{dJ}{d\gamma_p} = \frac{dJ}{dR} \frac{dR}{d\gamma_p} = 0.54 \gamma_p (\gamma_p^2 - 1)^{-1.625} \text{ protons/cm}^2\text{-sec-sr-unit } \gamma.$$

The computed source density using this spectrum in its restricted region of validity $2 \lesssim \gamma_p \lesssim 10^2$, is nearly the same numerically as that of equation (6).

The source density used by Brunstein (1965) for the energy interval $5 < \gamma_e < 50$ was $Q'(E_e) = 0.91 \rho E_e^{-2.625}$ electrons/cm³-sec-MeV, which can be transformed:

$$Q'(\gamma_e) = Q'(E_e) \frac{dE}{d\gamma} = 0.91 \rho (\gamma_e - 1)^{-2.625} (\text{mc}^2)^{-1.625} =$$

$$2.7 \rho (\gamma_e - 1)^{-2.625} \text{ electrons/cm}^3\text{-sec-unit } \gamma.$$

This source density, in its restricted region of validity, is also very close to that of equation (6).

b. Neutron-Decay Electrons

Cosmic rays produce secondary relativistic electrons also by means of nuclear reactions. In this process intermediate excited nuclei or neutrons are produced which in turn beta-decay. This two-step process systematically shifts the kinetic energies involved from the BeV region to the MeV region by giving the decay electrons a distribution in energy per unit mass resembling that of the parent neutrons or excited nuclei. Since the lifetime of the neutron against decay (1000 sec.) is infinitesimal compared to its lifetime against interaction, each secondary neutron will give rise to one secondary electron.

There are at least three sorts of cosmic-ray interactions with the interstellar medium which produce secondary neutrons. (1) Pi^+ mesons are produced with neutrons in the reaction $p + p \rightarrow p + n + \pi^+$. This process has the lowest energy threshold of any meson-producing reaction and excess π^+ production continues to occur in the higher-energy region of multiple π^+ and π^- meson production. Except for the case of the relatively rare process $p + p \rightarrow d + \pi^+$ which competes for π^+ production, charge conservation insures that the total number of excess π^+ mesons, integrated over all energies, is equal to the number of beta-decay electrons at production. (2) Neutrons also are made by the evaporation of excited compound nuclei formed in cosmic-ray interactions. Since about 10% of interstellar material may be helium and about 15% of cosmic rays are alphas or heavier nuclei, nearly one quarter of all cosmic-ray interactions with the interstellar medium involve compound nuclei capable of excitation. Neutron emission is the most likely method of the deexcitation of compound

nuclei. Although the neutrons are thermal in the reference frame of the nucleus, the cosmic-ray motion insures a spectrum up to relativistic energies in the observer's frame of reference. (3) Neutrons can also be made by a nuclear knock-on process analogous to that for atomic knock-on electrons. Neutrons made in this way have somewhat greater kinetic energies than do evaporation neutrons since less energy goes to the remaining nucleus.

The existence of these three processes requires that a secondary component of electrons should arise from nuclear reactions and also requires that this component should be characterized by energies in the MeV region. The spectrum of beta-decay electrons resulting from interstellar interactions can be reasonably estimated, but unlike that of knock-ons, cannot be calculated accurately since exact expressions for the nuclear reaction cross sections do not exist.

The probability of a neutron at rest emitting an electron within $d\gamma_1$ at γ_1 in its decay reaction $n \rightarrow e^- + p + \bar{\nu}$ is

$$f_1(\gamma_1) d\gamma_1 = 0.614 \gamma_1 (\gamma_1^2 - 1)^{1/2} (2.53 - \gamma_1)^2 d\gamma_1 \text{ electron/neutron.}$$

This beta-decay spectrum covers the kinetic-energy region from 0 to 782 keV, i. e., from a γ_1 of 1. to 2.53. The spectrum of electrons in the observer's frame, $f_3(\gamma_3)$ depends on both the spectrum of electrons relative to the parent neutrons $f_1(\gamma_1)$, and the spectrum of neutrons relative to the observer $f_2(\gamma_2)$.

Analogous to the calculation by Jones (1963) for $\pi \rightarrow \mu \rightarrow e$ electron productions, we have, for cases in which the distribution $f_3(\gamma_3)$ of electron energies in the decay frame does not depend on the spectrum $f_2(\gamma_2)$ of decay reactions relative to the observer, and for cases of isotropic collisions or decays,

$$f_3(\gamma_3) d\gamma_3 = \frac{1}{2} d\gamma_3 \int_1^{2.53} \frac{f_1(\gamma_1) d\gamma_1}{(\gamma_1^2 - 1)^{1/2}} \int_{\gamma_2^-}^{\gamma_2^+} \frac{f_2(\gamma_2) d\gamma_2}{(\gamma_2^2 - 1)^{1/2}} \quad (7)$$

Electrons produced in $d\gamma_1$ at γ_1 are uniformly spread over the interval

$$\gamma_2 \gamma_1 - (\gamma_2^2 - 1)^{1/2} (\gamma_1^2 - 1)^{1/2} \leq \gamma \leq \gamma_2 \gamma_1 + (\gamma_2^2 - 1)^{1/2} (\gamma_1^2 - 1)^{1/2}$$

(Rossi, 1954). Thus, the contribution to the interval $d\gamma_3$ at γ_3 from $d\gamma_1$ at γ_1 is integrated over the portion of the spectrum $f_1(\gamma_1)$ which contributed to $d\gamma_3$, using the connection between γ_1 and γ_3 expressed by the limits of integration on γ_2 :

$$\gamma_2^\pm = \gamma_1 \gamma_3 \pm (\gamma_1^2 - 1)^{1/2} (\gamma_3^2 - 1)^{1/2}$$

A neutron spectrum $f_2(\gamma_2)$ can be replaced by a discrete series of intensities of monoenergetic neutrons in order to obtain a numerical result. The

beta-decay spectra resulting from several monoenergetic neutron distributions are illustrated in Figure 4. The area $\int_1^{\infty} f_3(\gamma_3) d\gamma_3$ under each curve is 1.

The decay spectrum of the electron is given by equation (5), but the neutron spectrum relative to the galaxy $f_2(\gamma_2)$ depends, in turn, on both the neutron spectrum relative to the parent interaction $f_4(\gamma_4)$ and the spectrum of cosmic-ray interaction, relative to the galaxy $f_5(\gamma_5)$. The secondary neutron spectrum $f_4(\gamma_4)$ is a combination of at least three kinds of production spectra, as outlined earlier, and is unknown. However, it can be assumed that in the interaction, or center of momentum, frame, most of the secondary neutrons have energies in the few-MeV region, low compared with BeV-like cosmic-ray energies in the galactic frame. As shown by Jones (1963), a primary power-law spectrum, when such an inequality is obeyed, produces again a power-law spectrum of secondary products with the same shape but a different normalization. In the absence of detailed knowledge of the neutron production by cosmic rays, we can therefore assume that the neutron spectrum in the galactic frame reflects the cosmic ray spectrum with a multiplying constant containing the cross section for neutron production. We further simplify by using an energy-independent cross section of 10 millibarns, based on inspection of the data tabulated by Pollack and Fazio (1965). This appears to approximate the cross section for excess π^+ over π^- production, and, as discussed earlier, we can approximate by assuming that each excess π^+ is accompanied by one neutron. The source density of secondary neutron-decay electrons due to interactions of primary cosmic-ray protons is thus

$$Q_1(\gamma_3, \vec{r}) d\gamma_3 = 4\pi\rho(\vec{r}) f_3(\gamma_3) d\gamma_3 \text{ electrons/cm}^3\text{-sec,}$$

in which $f_3(\gamma_3)$ is related to $f_2(\gamma_2)$ by equation (7) and we approximate

$$f_2(\gamma_2) d\gamma_2 \approx N_o \sigma \frac{dJ_2}{d\gamma} d\gamma_2 \text{ electrons/sec-sr-gram.}$$

Here, $N_o = 6.02 \times 10^{23}$ per gram, $\sigma \approx 1.0 \times 10^{-26}$ cm² electron/proton and $dJ_2/d\gamma_2$ is the cosmic-ray spectrum of protons/cm²-sec-sr-unit γ given by equation (4). This calculated electron source density is found to be spectrally at least one decade below the knock-on source density. The beta-decay source density is thus so much less than the knock-on source density that, although its calculation is much less accurate, we assume its contribution can be neglected in the calculations that follow.

c. Galactic Energy-Loss Mechanisms

Electrons produced by the various cosmic-ray processes in the galaxy must subsequently lose energy in the galactic medium before they are observed. Since the medium is composed of matter, fields and radiation and has a boundary, the energy-loss mechanisms include ionization, bremsstrahlung, synchrotron radiation, inverse Compton effect and leakage into metagalactic space. Some of these are of greater importance than others; several authors have considered these effects for the higher-energy electrons from the $\pi \rightarrow \mu \rightarrow e$ process (e.g., Gould and Burbidge (1965), Ginzburg and Syrovatskii (1964), Jones (1963), Hayakawa and Okuda (1962) and others) and we use their commonly accepted values of astrophysical parameters in order to best compare results.

Ionization loss in the interstellar medium can be taken to be energy-independent over the electron energy range of interest here; although the loss does increase slightly at the highest energies, other loss mechanisms produce a much greater effect in this higher energy region. Hayakawa and Kitao (1956) have calculated the energy loss as a fraction of degree of ionization of the medium; if we assume the galactic medium to be mostly hydrogen and to be about 90 per cent neutral, then we can use a value of $\left| \frac{dE}{ds} \right| \approx 5 \text{ Mev cm}^2/\text{gram}$. The energy loss rate is directly proportional to the interstellar matter density; so if we then assume a constant $\rho_{\text{halo}} = 2 \times 10^{-26} \text{ g/cm}^3$, we can write

$$-\left(\frac{d\gamma}{dt}\right)_{\text{ion}} = \frac{c\rho}{mc^2} \left| \frac{dE}{ds} \right| = 6.0 \times 10^{-15} \equiv k \text{ unit } \gamma/\text{sec.} \quad (8)$$

In the disk, the value is about 50 times as great. It is interesting to note that the rectilinear range in the halo of even a low-energy relativistic electron is $\approx 10^{25} \text{ cm}$, much greater than the galactic diameter; in the disk where $\rho \approx 10^{-24} \text{ g/cm}^3$, it is somewhat less than the diameter.

Bremsstrahlung is the radiation which occurs when electrons decelerate in matter; the loss rate is proportional to the density and, using the values of the constants previously assumed, we can write

$$-\left(\frac{d\gamma}{dt}\right)_{\text{brem}} = \frac{4c r_e^2 Z_T^2 \rho N_o}{137 A_T} \ln\left(\frac{183}{Z_T^{1/3}}\right) \cdot (\gamma - 1) \approx 4.35 \times 10^{-18} \gamma \text{ unit } \gamma/\text{sec.}$$

This loss rate increases with energy, exceeding the rate due to ionization loss at energies above 700 MeV.

Synchrotron radiation, which occurs when electrons accelerate moving through magnetic fields, is an energy-loss mechanism which is independent of the matter density ρ , but depends on the square of the galactic field component B_{\perp} perpendicular to the electron's velocity. If we assume the mean B_{\perp} to be 3×10^{-6} gauss, we can write

$$-\left(\frac{d\gamma}{dt}\right)_{\text{syn}} = \frac{2cr_e^2 B_{\perp}^2 (\gamma^2 - 1)}{3mc^2} \simeq 1.73 \times 10^{-20} \gamma^2 \equiv b\gamma^2 \text{ unit } \gamma/\text{sec.} \quad (9)$$

In this case the rate increases by approximately the square of the energy, exceeding the rate due to ionization loss at energies above 300 MeV. Since the energy loss due to synchrotron radiation is greater than that due to bremsstrahlung at all energies for which either is more important than ionization loss, we will ignore the effects of bremsstrahlung in our calculation.

The energy loss due to inverse Compton collisions of electrons with interstellar photons is proportional to the radiation density and, in the region of interest here, to the square of the energy. If we assume that the typical galactic halo radiation density ρ_r is approximately

$$0.1 \text{ eV/cm}^3 = 1.6 \times 10^{-13} \text{ erg/cm}^3 ,$$

we can write

$$-\left(\frac{d\gamma}{dt}\right)_{\text{comp}} = \frac{-\rho_r}{B_{\perp}^2/4\pi} \left(\frac{d\gamma}{dt}\right)_{\text{syn}} \approx 3.9 \times 10^{-21} \gamma^2 \text{ unit } \gamma/\text{sec.} \quad (10)$$

This loss rate is less than, but has the same energy dependence as, the synchrotron rate over all the energies of interest here, $1 < \gamma \lesssim 10^5$. We shall add the contributions from the two effects in the work that follows.

The escape rate due to the diffusion of the electrons into metagalactic space is even less certain than the loss rates considered above, since the galactic trapping of electrons in the energy range of interest here is unknown. Electrons of these very low rigidities may be trapped in the galaxy for a time similar to the lifetime usually assumed to be representative of the high-energy cosmic-ray protons, or they may be trapped for a duration orders of magnitude longer. High-energy electrons are usually assumed to have the same lifetime against escape as do cosmic rays (Hayakawa and Okuda, 1962; Ginzburg and Syrovatskii, 1964; Gould and Burbidge, 1965), namely, $T_L = 3 \times 10^{15}$ sec. We will leave this an open question, and assume T_L for these low energies to be energy independent and to be between 3×10^{15} sec. and ∞ . For purposes of comparison, an effective energy loss rate is taken to be the total energy of the particle divided by the lifetime:

$$-\left(\frac{d\gamma}{dt}\right)_L = \frac{\gamma}{T_L} \leq 3.3 \times 10^{-16} \gamma \text{ unit } \gamma/\text{sec.}$$

The loss rate increases with energy, in the limit exceeding that rate due to ionization loss at energies above 9 MeV, and being exceeded by that due to synchrotron loss above 10 BeV. It thus may dominate ionization loss and synchrotron and other losses over most of the energy range of interest in this paper. This process, for purposes of computation, produces a particle sink, rather than an alteration of energy, and we therefore use

$$- Q_L(\gamma) = \frac{n(\gamma)}{T_L} \leq 3.3 \times 10^{-16} \cdot n \equiv \lambda n \text{ electrons/cm}^3\text{-sec-unit } \gamma .$$

d. Calculation of Secondary Galactic Electron Intensity

The intensity of low-energy electrons produced in the galaxy by cosmic rays and subject to the energy losses outlined above is calculated by use of the continuity equation for the particle density in γ space:

$$\frac{\partial n}{\partial t} + \text{div}_\gamma (n\dot{\gamma}) = S$$

Here $n(\gamma, t)d\gamma$ is the spatial density in $d\gamma$ at γ of electrons/cm³ and $S(\gamma)d\gamma$ the source production rate of electrons/cm³-sec. The continuity equation is rewritten then, as

$$\frac{\partial n}{\partial t} + \frac{\partial}{\partial \gamma} n \left(\sum_i \frac{d\gamma_i}{dt} \right) = \sum_i Q_i$$

in which the right hand side is the algebraic sum of the particle sources and sinks Q_i , and

$$\frac{\partial}{\partial \gamma} \left(n \sum_i \frac{d\gamma_i}{dt} \right) d\gamma dt$$

is the net increase in the spatial density of electrons in $d\gamma$ at γ due to the various energy loss or gain mechanisms. Inserting the production and escape terms Q_i and the energy loss terms $\frac{d\gamma_i}{dt}$ discussed earlier, we have

$$\frac{\partial n}{\partial t} - (k + b\gamma^2) \frac{\partial n}{\partial \gamma} + (\lambda - 2b\gamma) n = Q_1(\gamma) \quad (11)$$

in which k , b and λ represent the positive numerical coefficients outlined above for ionization loss, synchrotron loss, and escape, respectively, and Q_1 refers to the knock-on source density. The solution to this equation for $n(\gamma)$ is shown in the Appendix to have various forms, depending on which various energy-loss mechanisms are taken to be dominant. Since knowledge of the astrophysical parameters is uncertain, we exhibit here all the solutions.

First, in the limiting case at the lowest energies when only ionization losses and escape losses are relevant,

$$n_e(\gamma) \Big|_{b=0} = \frac{e^{\lambda\gamma/k}}{k} \int_{\gamma}^{\gamma+tk} Q(\gamma') e^{-\lambda\gamma'/k} d\gamma' \text{ electrons/cm}^3\text{-unit } \gamma.$$

If we use for $Q(\gamma')$ the knock-on source expression of equation (6), and take $t \rightarrow \infty$, the integral can be analytically evaluated for $\lambda \rightarrow 0$, and can be numerically evaluated for finite λ . The resulting differential intensities

$$\frac{dj_e}{d\gamma} = \frac{cn_e}{4\pi} \text{ electrons/cm}^2\text{-sec-sr-unit } \gamma$$

are plotted in Figure 5.

Second, in the much higher-energy region when synchrotron losses predominate,

$$\frac{dj_e}{d\gamma} = \frac{ce^{f(k,\lambda,b,\gamma)}}{4\pi(k+b\gamma^2)} \int_{\gamma}^{\infty} Q(\gamma') e^{-f(k,\lambda,b,\gamma')} d\gamma' \text{ electrons/cm}^2\text{-sec-sr-unit } \gamma,$$

in which

$$f(k, \lambda, b, \gamma) = \frac{\lambda}{(kb)^{1/2}} \arctan \frac{b^{1/2}\gamma}{k^{1/2}}$$

In the event of no escape losses, we have

$$\left. \frac{d j_e}{d \gamma} \right|_{\lambda = 0} = \frac{c}{4 \pi (k + b \gamma^2)} \int_{\gamma}^{\infty} Q(\gamma') d \gamma'$$

These calculated intensities are also plotted in Figure 5. Comparison of the calculated secondary intensities with the intensity observed by Cline, et al. (1964) shows that the calculated absolute intensity is too low. Further, using the modulation model of Parker (1963), the observed value would be a factor of 2.7 higher, increasing the discrepancy. We claim that this discrepancy is not trivial; i. e., the lack of knowledge of astrophysical parameters is not a contributing factor in this lack of fit. In the limiting case in which only ionization losses contribute, the secondary electrons are in an equilibrium state such that the matter density cancels out:

$$\left. \frac{d j_e}{d \gamma} \right|_{\substack{b = 0 \\ \lambda = 0}} = \frac{c}{4 \pi k} \int_{\gamma}^{\infty} Q(\gamma') d \gamma' = \frac{c}{4 \pi k} \left(\frac{4.9 \rho}{1.76 (\gamma - 1)^{1.76}} \right)$$

electrons/cm²-sec.-sr-unit γ , in which k is directly proportional to ρ . Thus, since the ionization losses and knock-on cross section are well known, the calculated intensity in this case depends only upon the assumption of the galactic universality of the cosmic-ray intensity.

III. SECONDARY METAGALACTIC ELECTRONS

a. Production Mechanisms

Secondary electrons can be made in metagalactic space by cosmic-ray interactions with that medium. Although the character of the intergalactic cosmic-ray intensity and of the medium are poorly known, we can still expect that the secondary low-energy electrons will be in an equilibrium similar to that in the galaxy. As was seen in the previous section, the fundamental quantity to know in order to calculate the secondaries is the primary cosmic-ray intensity. Some theorists assume that the metagalactic intensity is about the same as that observed in the galaxy; for purposes of calculation we will take it to be an upper limit.

Although the degree of ionization of the intergalactic medium is uncertain, we can make the usual assumption that it is essentially totally ionized; the cross section for knock-on production by Coulomb collisions with free plasma electrons is essentially the same as that for bound electrons. Nuclear interactions of cosmic rays with free protons will produce neutron-decay electrons as well, but again to a lesser extent; in fact, evaporation neutron production will be even less competitive in intergalactic space if we assume that both the cosmic rays and the medium contain negligible portions of alpha particles.

We then take the metagalactic secondary source density to be given by

$$Q_m(\gamma_e) \lesssim \frac{4.9}{1.75} \rho (\gamma_e - 1)^{-2.76} \approx 2.8 \rho (\gamma_e - 1)^{-2.76} \text{ electrons/cm}^3\text{-sec-unit } \gamma,$$

in which ρ is the density in g/cm^3 .

b. Metagalactic Energy-Loss Mechanisms

Secondary electrons produced in intergalactic space lose energy because of their interaction with that medium, and also vanish because of the universal expansion. Assuming knowledge of the physical parameters involved, these losses can be calculated. In addition, there may be a distortion of the energy spectrum due to the effects of penetration into the galaxy, just as there is into the interplanetary region.

Ionization losses in a fully ionized medium are replaced by losses due to plasma oscillations; the cosmic-ray electron loses energy by interacting with the plasma ions and electrons to a greater extent than it would in a neutral medium since the shielding of the individual charges is reduced. As shown by Hayakawa and Kitao (1956), for an ionized medium we have

$$\left| \frac{dE}{ds} \right|_{mg} = 10 \text{ Mev-cm}^2/\text{gram.}$$

If we assume a universal proton density of $10^{-5}/\text{cm}^3$, or an equivalent matter density $\rho_{mg} = 1.6 \times 10^{-29} \text{ g/cm}^3$, using equation (8) we have for the energy loss rate

$$-\left(\frac{d\gamma_e}{dt} \right)_{ion} = 1.0 \times 10^{-17} \text{ unit } \gamma/\text{sec.}$$

It is interesting to note that the rectilinear range of a low-energy relativistic electron is about 10^{28} cm, close to the Hubble radius. Thus, electrons produced universally can reach the galaxy. The ratio of metagalactic to galactic bremsstrahlung losses is the same as that ratio for "ionization" losses. This is because both interactions depend linearly on the matter density, and because a factor of 2 is introduced in each loss when one substitutes an ionized medium for a nearly neutral one. Thus

$$-\left(\frac{d\gamma_e}{dt}\right)_{\text{brem}} \cong 7.0 \times 10^{-21} \gamma_e \text{ unit } \gamma/\text{sec.}$$

The bremsstrahlung loss equals that due to plasma excitation at $\gamma_e = 1.4 \times 10^3$ or about 700 MeV.

Synchrotron and inverse Compton losses depend on the typical metagalactic magnetic field strength and photon density, which are unknown. Gould and Burbidge (1965) adopt 10^{-7} gauss, and $6.0 \times 10^{-3} \text{ eV/cm}^3 \approx 10^{-14} \text{ erg/cm}^3$ respectively, as typical assumptions. For purposes of comparison, then, we have from equations (9) and (10)

$$-\left(\frac{d\gamma_e}{dt}\right)_{\text{syn}} \cong 2.0 \times 10^{-23} \gamma^2 \text{ unit } \gamma/\text{sec}$$

and

$$-\left(\frac{d\gamma_e}{dt}\right)_{\text{comp}} \approx 2.4 \times 10^{-22} \gamma^2 \text{ unit } \gamma/\text{sec}.$$

Inverse Compton losses thus dominate synchrotron losses at all energies. Since plasma-excitation losses dominate inverse Compton losses for all energies up to 100 MeV, well above the energy region where bremsstrahlung competes, Compton effect is the only high-energy loss mechanism we will consider.

The magnitude of the effective particle sink represented by universal expansion, the red-shift loss, is uncertain since the diffusion of electrons in intergalactic space cannot be accurately represented. We will therefore, analogous to the galactic case, take as an upper limit the rectilinear loss due to the three-dimensional expansion, characterized by a lifetime one-third the Hubble factor: $T_L = 1/3H$. The equivalent energy loss rate is

$$-\left(\frac{d\gamma_e}{dt}\right)_L \leq \frac{\gamma}{T_L} \leq 1.0 \times 10^{-17} \gamma \text{ unit } \gamma/\text{sec}.$$

This loss rate is exceeded by that due to inverse Compton effect above 20 BeV but it actually exceeds that due to plasma excitation at all energies above $\gamma = 1$.

For our computation we represent **this process** as a sink of particles and so

$$- Q_L(\gamma) = \frac{n(\gamma)}{T_L} \leq 1.0 \times 10^{-17} \cdot n \text{ electrons/cm}^3\text{-sec-unit } \gamma.$$

c. Calculation of Secondary Metagalactic Electron Intensity

The intensity of low-energy electrons produced in intergalactic space by cosmic rays with an assumed universal intensity is calculated by use of the equations derived earlier. Using the values stated above for intergalactic plasma-excitation and inverse Compton losses, we solve for two extreme cases, namely, that for the rectilinear red-shift loss and that for none. These are plotted in Figure 5. It is seen that the metagalactic flux, even for a primary cosmic-ray flux equal to that observed here, is at least an order of magnitude below the calculated galactic electron flux. The analogous result for the high-energy secondaries was obtained by Gould and Burbidge (1965). Thus, since this calculated metagalactic electron intensity is an upper limit, it represents a negligible addition to the galactic secondary flux, and we must look elsewhere for a possible source for the experimentally observed electrons.

IV. ENERGY INPUTS TO SECONDARY ELECTRONS

In this section we consider the possibility that the observed low-energy electrons are galactic secondaries which have been accelerated after their

creation. Two mechanisms for increasing the electron intensity to that observed present themselves, one in the solar system producing a local intensity increase, and the other producing an intensity increase throughout interstellar space.

a. Heliocentric Electric Field

The possibility of the presence of a substantial potential of the solar system with respect to the galaxy as a whole has been considered by several authors. In particular, attempts have been made to explain the solar modulating effects of primary cosmic-rays in terms of such a potential. Earlier work by Ehmert (1960) required a positive potential of 1 or more BV to account for the observed modulation. A more recent treatment by Freier and Waddington (1965) required 50 MV to obtain quantitative agreement between their model and the observed spectrum. It is nevertheless difficult to envision how even such a modest electric field could be maintained in the presence of the interstellar gas which is, undoubtedly, somewhat ionized.

In this section, we consider the effects of such a heliocentric electric field on the interstellar secondary electrons, which we have shown must be produced with an intensity not far from that observed.

The Liouville theorem states that the differential flux of electrons

$$\frac{dJ}{(4\pi r^2 dr)(4\pi p^2 dp)}$$

remains constant in phase space. From this, we find that

$$\left(\frac{dJ}{d\gamma}\right) \cdot \left(\frac{1}{p^2}\right)$$

is a constant. Relating the spectrum outside the solar system, $\frac{dJ_1}{d\gamma}$, to that observed, $\frac{dJ_2}{d\gamma}$, we have

$$\frac{dJ_2}{d\gamma} \cdot \left\{(\gamma mc^2 + qV)^2 - (mc^2)^2\right\}^{-1} = \frac{dJ_1}{d\gamma} \left\{(\gamma mc^2)^2 - (mc^2)^2\right\}^{-1}$$

where qV is the product of the charge on the electrons and the accelerating potential. Relating the measured intensity, defined here as $\frac{dJ_2}{d\gamma} = \frac{dJ}{d\gamma} (\gamma mc^2 + qV)$, to the calculated secondary intensity, $\frac{dJ_1}{d\gamma} = \frac{dJ}{d\gamma} (\gamma mc^2)$, we find that a potential of 1.5 million volts provides an excellent fit to the data at the low energies. This result is shown in Figure 6.

The intensity increase provided by this 1.5 MV field is insufficient to simultaneously fit the observations at the high energies. The negative electrons in the fractional-BeV energy region, found to be in excess of those from $\pi \rightarrow \mu \rightarrow e$ decay by DeShong, Hildebrand, and Meyer (1964), would thus have to be due to yet another source. Such may or may not be the case, but a third source is undesirable from the point of view of minimizing hypotheses. The observations may yet indicate that there could be a change in the spectral form between 20 and

200 MeV, but it will have to be determined whether this is due to solar modulation. If we assume that there is only one electron source other than the $\pi \rightarrow \mu \rightarrow e$ process, we thus need to look elsewhere for the required energy input. Further, a 50 MV field would so grossly distort the 3-MeV electron spectrum from galactic knock-on production as to be experimentally evident from a vast over-supply of low-energy electrons.

b. Galactic Fermi Acceleration

The acceleration mechanism of cosmic-ray protons most generally considered as a possible means of interstellar energy increase is the mechanism first investigated by Fermi (1949). In this model, a particle randomly gains energy in the process of colliding with moving magnetic irregularities in the medium in which it is confined. The probability for it to have total energy between γ and $\gamma + d\gamma$ is

$$\frac{d\pi}{d\gamma}(\gamma) d\gamma = \frac{1}{T\alpha} \frac{\gamma_0^{1/T\alpha}}{\gamma^{1+1/T\alpha}} d\gamma ,$$

in which γ_0 is the particle's value at injection, T is the mean life of the particles, and $\alpha \equiv v^2/(c^2\tau)$, in which v is the scattering center velocity and τ is the mean time between collisions. Thus,

$$\frac{dJ}{d\gamma} = \frac{J_0}{T\alpha} \frac{\gamma_0^{1/T\alpha}}{\gamma^{1+1/T\alpha}}$$

in which

$$J_0 \equiv \int_1^{\infty} \frac{dJ}{d\gamma} d\gamma ,$$

showing that the resultant cosmic-ray spectrum should obey a power law in total energy. Since this result is, in fact, consistent with observations, a mechanism of the Fermi type gains much support, whether operative in the interstellar medium, or in stellar atmospheres, or in supernovae.

If we now investigate the possibility that electrons may also be Fermi accelerated, we treat the mechanism as an energy input in the solution of the continuity equation for electrons in energy space. Since, in the Fermi model, the energy acquired by a particle of age t is $\gamma = e^{at}$ in which a is defined above, we have

$$+ \left(\frac{d\gamma}{dt} \right)_{\text{Fermi}} = a \cdot \gamma \text{ unit } \gamma/\text{sec.}$$

Inserting this term in the continuity equation, we have

$$\frac{\partial n}{\partial t} - (k - a\gamma + b\gamma^2) \frac{\partial n}{\partial \gamma} + (a + 2b\gamma + \lambda)n = Q(\gamma) \quad (12)$$

in place of equation (11). Since there now exist both positive and negative energy loss terms, the solutions to this equation do not directly follow from those used earlier. As shown in the Appendix, the forms of the solutions vary, depending on the choice of the parameters k , α , b and λ . In particular, the algebraic sign of $(\alpha^2 - 4kb) \equiv \nu^2$ is the determining factor; i. e., whether α is greater or smaller than about $2 \times 10^{-17} \text{ sec}^{-1}$, using a halo density of $2 \times 10^{-26} \text{ g cm}^{-3}$. As we shall see, any value of α that comes close to giving a fit to the experimental data is much greater than this; so we can, except for the postulated $\alpha \equiv 0$ case discussed earlier, take $(\alpha^2 - 4kb)$ to be positive.

As discussed in the Appendix, care must be taken that the behavior is correctly interpreted near the singularities which occur at energies

$$\gamma^{\pm} \equiv \frac{1}{2b} \left\{ \alpha \pm (\alpha^2 - 4kb)^{\frac{1}{2}} \right\}$$

where the rate of energy loss changes signs. The solutions for

$$\frac{dj_e}{d\gamma} = \frac{cn_e}{4\pi}$$

are described for the various cases of escape and synchrotron losses being or not being incorporated. For $\gamma < \gamma^+$, which we shall see is the energy range of

interest here, the general solution is

$$\frac{dj_e}{d\gamma} = \frac{c}{4\pi(k - a\gamma + b\gamma^2)} \left(\frac{\gamma^+ - \gamma}{\gamma - \gamma^-} \right)^{\lambda/\nu} \int_{\gamma}^{\gamma^-} Q(\gamma') \left(\frac{\gamma' - \gamma^-}{\gamma^+ - \gamma'} \right)^{\lambda/\nu} d\gamma'.$$

Its limiting form for $\lambda \rightarrow 0$ follows directly but for $b \rightarrow 0$ it is more convenient to solve the original differential equation to yield

$$\left. \frac{dj_e}{d\lambda} \right|_{b=0} = \frac{c}{4\pi k (1 - a\gamma/k)^{1 + \lambda/a}} \int_{\gamma}^{k/a} Q(\gamma') (1 - a\gamma'/k)^{\lambda/a} d\gamma'$$

which also has an evident limit for $\lambda \rightarrow 0$. These results are plotted in Figure 7 for various values of α . If α is large enough it is possible to match the intensity but with a poor fit to the spectral shape.

If one attributes the primary proton beam to Fermi acceleration and the further assumption is made that the parameters α and λ are nearly the same for protons and electrons, as shown by Fermi (1949), we have for the proton beam

$$\frac{dJ}{d\gamma} = \text{const. } \gamma^{-(1+\lambda/a)}.$$

The exponent $(1 + \lambda/\alpha)$ is well established to be approximately 2.5. We now must have $\lambda > \alpha$ instead of $\lambda \lesssim \alpha$. Here $\lambda = 1.5\alpha$, and the singularity in the differential electron spectrum is removed. This solution is shown in Figure 8 for a variety of values of λ and α , including those used formerly. In this case, toward the lower energies the resulting spectrum becomes independent of λ and α and has the same slope as the observed spectrum, but, like the solution for $\alpha = 0$ which it approaches, is too low in intensity. It appears that if the electrons are propagated similarly to the protons, such that a Fermi acceleration with $\lambda/\alpha = 1.5$ is present, the secondaries from interstellar knock-on production cannot account for the observations. Of course, it is not obvious that either λ or α would have the same values for electrons as they have for protons, or that λ/α must necessarily be equal to 1.5. The electrons have a much lower rigidity than the protons do at a given γ ; they would therefore have a greater lifetime against escape and possibly a much different acceleration.

We can conclude that the intensity and shape of the observed low-energy electron spectrum cannot simultaneously be accounted for by galactic knock-on secondaries.

V. CONCLUSION

The secondary electron spectrum to be expected from $\pi \rightarrow \mu \rightarrow e$ decay as calculated by Hayakawa and Okuda (1962), Jones (1963), and others, coupled with the positron-electron measurements by DeShong, et al. (1964) and by Hartman, et al. (1965), indicate that meson-decay electrons cannot adequately explain the observed properties of high-energy cosmic-ray electrons.

Two of the authors have in a previous paper (Brunstein and Cline [1966]) discussed the possibility that the electrons may be of a primary nature, being accelerated in the same process along with cosmic-ray protons. We have here examined the possibility that cosmic-ray electrons are secondary to the nuclear beam, arising from both knock-on collisions and from the beta-decay of the resultant neutrons from nuclear interactions in the interstellar gas and in metagalactic space. We conclude that neutron-decay electrons are well below the intensity to be expected from knock-on collisions and thus may be neglected as a potential source. However, at least at low energies, an intrinsically accurate calculation of the knock-on electron intensity shows it to be well above that expected from $\pi \rightarrow \mu \rightarrow e$ decay but substantially below the measured intensity of low-energy electrons.

We have therefore postulated two prospective models to account for the observed electrons as being due to these knock-on electrons: (1) the knock-on electrons are further accelerated after their injection by a heliocentric electric field, (2) the knock-on electrons are accelerated after injection by a general galactic Fermi mechanism.

The first possibility, the heliocentric electric field, requires only a modest 1.5 MV to boost the calculated knock ons to the observations; however, it does not appreciably affect the higher-energy electrons where we still must face the question of the excessive negative electrons. Moreover, it is difficult to envision how even a modest potential of 1.5 MV could be maintained by the solar system (Lam and Sandri [1965]).

The second possibility, that of Fermi acceleration after secondary production, is seen to have limitations. The value of the parameter α , defined by

$\frac{d\gamma}{dt} = \alpha\gamma$, is varied in the neighborhood of that value which produces an approximate fit to the data, $\alpha \approx 10^{-15} \text{ sec}^{-1}$. When a lifetime against escape of about 10^8 years is assumed ($\lambda \equiv 3.3 \cdot 10^{-16} \text{ sec}^{-1}$) the observed intensity can be roughly fitted, although the fit to the slope in the low-energy region is poor. In the high-energy region, if we assume that about half of the electrons are accounted for by $\pi^- \mu^- e$ decay, we find that the remainder can also be roughly fitted. However, in this case, the value of α is quite large, $> 10^{-15} \text{ sec.}$, and is about the same as that value shown by Brunstein and Cline (1966) to describe the electron intensity in terms of Fermi-accelerated primary thermal electrons. If these parameters accurately represent galactic conditions, then we may draw the conclusion that the cosmic-ray beam will be composed of both secondary knock-on electrons and primary electrons Fermi-accelerated in essentially equivalent fractions. If the actual value of the parameter α is smaller, the secondary electrons account for a smaller fraction of the observed intensity. On the other hand, if the value of the ratio λ/α is a constant 1.5 as is usually assumed for protons, then in the low-energy region a fit to the observations, independent of α , is impossible.

We can conclude, in any case, that the secondary hypothesis alone cannot account for the observations and that at least half of the observed low-energy electrons have a primary origin. We note that the observations of the charge ratio led to a similar conclusion for the high-energy electrons (DeShong et al., 1964, Hartman, et al., 1965). We feel, therefore, that a primary cosmic-ray beam of electrons is present throughout the energy region from $\gamma \gtrsim 1$ to $\gamma > 10^5$, having about the same velocity spectrum as the cosmic-ray protons in the $1 < \gamma < 10^2$ region. However, further observations in the $\gamma < 5$ and $\gamma > 10^5$

region are desirable to better understand the properties of these primaries,
and more details in the $20 < \gamma < 200$ region are desirable to map the transition
between the knock-on and the $\pi^- \mu^- e$ secondary contributions.

APPENDIX

The equation to be solved is

$$\frac{\partial n}{\partial t} - (k - \alpha\gamma + b\gamma^2) \frac{\partial n}{\partial \gamma} + (a - 2b\gamma + \lambda)n = Q(\gamma), \quad (1)$$

in which $n = n(\gamma, t)$. The initial condition assumed is $n(\gamma, t = 0) = 0$, and the solution is obtained by the method of characteristics.

Before proceeding to the actual solution we consider two possibilities:

1. Particles produced with certain energies lose some of their energy when propagating through the medium, while others, produced with energies in a different range, gain energy from the medium. This implies that $\frac{d\gamma}{dt}$ changes sign, and for this to be so it is necessary that

$$\nu^2 \equiv a^2 - 4kb \quad (2)$$

be greater than zero. The changes in sign occur at the energies

$$\gamma^\pm = \frac{a \pm \nu}{2b}. \quad (3)$$

When $4kb/\alpha^2 \ll 1$, we have the approximations

$$\left\{ \begin{array}{l} \gamma^+ \approx \frac{a}{b} \\ \gamma^- \approx \frac{k}{a} \end{array} \right\}$$

2. Particles produced with any energy never gain energy from the medium.

This implies that for all energies $\frac{d\gamma}{dt} < 0$, and hence that

$$\mu^2 \equiv 4kb - \alpha^2 \tag{4}$$

be greater than zero.

We treat these two cases separately.

a. Predominance of Energy Increase

Here $\alpha^2 - 4kb > 0$.

The solution for this case is

$$n(\gamma, t) = \frac{(\gamma - \gamma^+)^{\lambda/\nu - 1}}{(\gamma - \gamma^-)^{\lambda/\nu + 1}} \int_{\gamma}^{g(\gamma, t)} Q(\gamma') \left(\frac{\gamma' - \gamma^-}{\gamma' - \gamma^+} \right)^{\lambda/\nu} d\gamma', \tag{5}$$

where

$$g(\gamma, t) = \frac{\gamma^+ (\gamma - \gamma^-) e^{-\nu t} - \gamma^- (\gamma - \gamma^-)}{(\gamma - \gamma^-) e^{-\nu t}} \quad (6)$$

At this stage we want to investigate the behavior of $N(\gamma, t)$ at the exceptional points $\gamma = \gamma^\pm$. One finds, by using the L'Hopital rule and the form (6) of $f(\gamma, t)$, that

$$\lim_{\gamma \rightarrow \gamma^-} n(\gamma, t) = \frac{Q(\gamma^-)}{\lambda + \nu} (1 - e^{-(\lambda + \nu)t}) . \quad (7)$$

Therefore at $\gamma = \gamma^-$, there is no singularity in $n(\gamma, t)$ for all times.

To find the limiting value of $n(\gamma, t)$ when $\gamma \rightarrow \gamma^+$, it is necessary to go to the original equation (1) and proceed there to the limit, since precisely at this energy there is a singularity also in the integrand of (5) in addition to the one outside the integral. Finally we obtain

$$\lim_{\gamma \rightarrow \gamma^+} n(\gamma, t) = \frac{Q(\gamma^+)}{\lambda - \nu} (1 - e^{-(\lambda - \nu)t}) . \quad (8)$$

Equation (8) shows that two possibilities may arise:

i) $\lambda - \nu > 0$:

Here there is no singularity at $\gamma = \gamma^+$ for all times, and

$$n(\gamma^+, t = \infty) = \frac{Q(\gamma^+)}{\lambda - \nu} \quad (9)$$

ii) $\lambda - \nu \leq 0$:

Here there is no singularity for finite times but when $t \rightarrow \infty$, then

$$n(\gamma^+, t = \infty) = \infty. \quad (10)$$

This situation persists when $\lambda \rightarrow 0$.

In order to determine, (for case ii), the behavior of $n(\gamma, t)$ when a steady state prevails, we must investigate the rate of energy loss,

$$\frac{d\gamma}{dt} = - (k - a\gamma + b\gamma^2) \quad (11)$$

and find the time intervals needed by particles produced at certain energies to achieve different energies. One finds that for all $\gamma < \gamma^+$, a steady state will be reached when $t \rightarrow \infty$; while for $\gamma > \gamma^+$, the steady state is attained after a finite time interval t_{\max} given by

$$t_{\max} = - \int_{\gamma}^{\infty} \frac{d\gamma'}{k - a\gamma' + b(\gamma')^2} = \frac{1}{\nu} \ln \frac{\gamma - \gamma^-}{\gamma - \gamma^+}. \quad (12)$$

One finds that when $t \rightarrow \infty$, $f(\gamma, t) \rightarrow \gamma^-$; and when $t \rightarrow t_{\max}$, $f(\gamma, t) \rightarrow \infty$.

Therefore the steady state solution $n_s(\gamma)$ is:

$$n_s(\gamma) = \begin{cases} \frac{1}{k - a\gamma + b\gamma^2} \left(\frac{\gamma^+ - \gamma}{\gamma - \gamma^-} \right)^{\lambda/\nu} \int_{\gamma}^{\gamma^-} Q(\gamma') \left(\frac{\gamma' - \gamma^-}{\gamma^+ - \gamma'} \right)^{\lambda/\nu} d\gamma', & \text{for } \gamma < \gamma^+ \\ \frac{1}{k - a\gamma + b\gamma^2} \left(\frac{\gamma - \gamma^+}{\gamma - \gamma^-} \right)^{\lambda/\nu} \int_{\gamma}^{\infty} Q(\gamma') \left(\frac{\gamma' - \gamma^-}{\gamma' - \gamma^+} \right)^{\lambda/\nu} d\gamma', & \text{for } \gamma > \gamma^+. \end{cases} \quad (13)$$

Note that for $\lambda > \nu$, $n_s(\gamma)$ has no singularities; while for $\lambda \leq \nu$, the only singularity is located at $\gamma = \gamma^+$, corresponding to a pile-up of particles on the energy axis where synchrotron loss equals Fermi-acceleration gain.

The integral appearing in (13) cannot be evaluated in closed form for arbitrary values of λ/ν . For the special case $\lambda \rightarrow 0$ we have

$$\lim_{\lambda \rightarrow 0} n_s(\gamma) \equiv n_s(\gamma) = \begin{cases} \frac{1}{k - a\gamma + b\gamma^2} \int_{\gamma}^{\gamma' \approx k/a} Q(\gamma') d\gamma', & \text{for } \gamma < \gamma^+ \\ \frac{1}{k - a\gamma + b\gamma^2} \int_{\gamma}^{\infty} Q(\gamma') d\gamma', & \text{for } \gamma > \gamma^+. \end{cases} \quad (14)$$

If $b = 0$, the equation for $n(\gamma, t)$ is

$$\frac{\partial n}{\partial t} - (k - a\gamma) \frac{\partial n}{\partial \gamma} + (a + \lambda) n = Q(\gamma) \quad (15)$$

and the solution is

$$n(\gamma, t) = \frac{1}{k(1 - a\gamma/k)^{1 + \lambda/a}} \int_{\gamma}^{f(\gamma, t)} Q(\gamma') \left(1 - \frac{a\gamma'}{k}\right)^{\lambda/a} d\gamma' \quad (16)$$

where

$$f(\gamma, t) \equiv \frac{k - (k - \alpha\gamma) e^{-\alpha t}}{\alpha}.$$

There is no singularity; in fact at the point $\gamma = \frac{k}{\alpha} \equiv \gamma_0$,
we obtain

$$\lim_{\gamma \rightarrow \gamma_0} n(\gamma, t) = \frac{1}{\alpha + \lambda} Q(\gamma_0) (1 - e^{-\alpha t}).$$

The steady state solution is obtained when we let $t \rightarrow \infty$:

$$n_s(\gamma) = \frac{1}{k(1 - \alpha\gamma/k)^{1 + \lambda/a}} \int_{\gamma}^{\gamma_0 \equiv k/\alpha} Q(\gamma') \left(1 - \frac{\alpha\gamma'}{k}\right)^{\lambda/a} d\gamma'$$

for $\gamma < \gamma_0$, and

$$n_s(\gamma) = \frac{1}{k(\alpha\gamma/k - 1)^{1+\lambda/a}} \int_{\gamma_0 \equiv k/a}^{\gamma} Q(\gamma') \left(\frac{\alpha\gamma'}{k} - 1\right)^{\lambda/a} d\gamma'$$

for $\gamma > \gamma_0$.

For $\lambda \rightarrow 0$ (or $\lambda \ll \alpha$), the steady state solution is:

$$n_s(\gamma) = \frac{1}{k - \alpha\gamma} \int_{\gamma}^{k/a} Q(\gamma') d\gamma' \quad (20)$$

which is the special case evaluated by Brunstein (1965).

b. Predominance of Energy Decrease

Here $\mu^2 \equiv 4kb - \alpha^2 > 0$.

The solution for this case is

$$n(\gamma, t) = \frac{\exp\left(\frac{2\mu}{\lambda} \arctan \frac{2b\gamma - \alpha}{\mu}\right)}{k - \alpha\gamma + b\gamma^2} \int_{\gamma}^{h(\gamma, t)} Q(\gamma') \exp\left(\frac{-2\lambda}{\mu} \arctan \frac{2b\gamma' - \alpha}{\mu}\right) d\gamma', \quad (21)$$

where

$$h(\gamma, t) = \frac{a}{2b} - \left(\frac{\mu}{2b}\right) \cdot \frac{a - 2b\gamma - \mu \tan(\mu t/2)}{\mu + (a - 2b\gamma) \tan(\mu t/2)}. \quad (22)$$

Equation (21) implies an electron density periodic in time, a result which obviously is inadmissible if not interpreted correctly. We show that beyond a certain characteristic time t_{\max} the expression (21) for $n(\gamma, t)$ loses all meaning. This follows immediately from equation (11), when we note that for the case under consideration

$$\frac{d\gamma}{dt} = -b \left\{ \left(\gamma - \frac{a}{2b} \right)^2 + \left(\frac{\mu}{2b} \right)^2 \right\} < 0. \quad (23)$$

The particles therefore always decelerate. We define t_{\max} as the time needed for a particle of infinite initial energy to decelerate to energy γ :

$$-t_{\max}(\gamma) = \frac{1}{b} \int_{\infty}^{\gamma} \frac{d\gamma'}{\left(\gamma' - \frac{a}{2b} \right)^2 + \left(\frac{\mu}{2b} \right)^2}, \quad (24)$$

or

$$t_{\max}(\gamma) = \frac{\pi}{\mu} - \frac{2}{\mu} \arctan \frac{2b\gamma - a}{\mu}. \quad (25)$$

It can be shown that at $t = t_{\max}$, $n(\gamma, t)$ reaches its first maximum. This follows from the fact that $Q(\gamma)$ is given by a negative power law and from the result that

$$\lim_{t \rightarrow t_{\max}} h(\gamma, t) = \infty. \quad (26)$$

This result would not be true for a production rate with a positive power law form.

Beyond t_{\max} , $n(\gamma, t)$ decreases toward zero. We therefore adopt the solution attained at $t = t_{\max}$ as the appropriate steady state solution. Thus

$$n_s(\gamma) = \frac{\exp\left(\frac{2\lambda}{\mu} \arctan \frac{2b\gamma - a}{\mu}\right)}{k - a\gamma + b\gamma^2} \int_{\gamma}^{\infty} Q(\gamma') \exp\left(\frac{-2\lambda}{\mu} \arctan \frac{2b\gamma' - a}{\mu}\right) d\gamma', \quad (27)$$

Again the integral in equation (27) cannot be evaluated in a closed form for arbitrary λ . For the case $\lambda \rightarrow 0$ (or $\lambda \ll \mu$), we can write

$$\lim_{\lambda \rightarrow 0} n_s(\gamma) \equiv n_s(\gamma) = \frac{1}{k - a\gamma + b\gamma^2} \int_{\gamma}^{\infty} Q(\gamma') d\gamma'. \quad (28)$$

Finally, the solution for $a \rightarrow 0$ follows from equation (27), or from (28) if $\lambda \rightarrow 0$, unless $b = 0$. In that case we again solve equation (1) and find

$$n(\gamma) \left| \begin{array}{l} \\ b = 0 \\ a = 0 \end{array} \right. = \frac{\exp(\lambda\gamma/k)}{k} \int_{\gamma}^{\infty} Q(\gamma') \exp\left(\frac{-\lambda\gamma'}{k}\right) d\gamma'. \quad (29)$$

REFERENCES

- Agrinier, B., Koechlin, Y., Parlier, B., Boella, G., Degli Antoni, G., Dilworth, C., Scarsi, L., and Sironi, G. 1964, Phys. Rev. Letters, 13, 377.
- Balasubrahmanyam, V. K., and McDonald, F. B. 1964, J. Geophys. Res., 69, 3289.
- Balasubrahmanyam, V. K., Boldt, E. A., and Palmeira, R. A. R. 1965, Phys. Rev., 140, B1157.
- Bhabha, H. J. 1938, Proc. Roy. Soc. (London), A164, 257.
- Burnstein, K. A. 1965, Phys. Rev., 137, B757.
- Burnstein, K. A., and Cline, T. L. 1966, Nature, to be published; also 1965, NASA, GSFC Energetic Particles Preprint X-611-65-479.
- Cline, T. L., Ludwig, G. H., and McDonald, F. B. 1964, Phys. Rev. Letters, 13, 786.
- Daniel, R. R., and Stephens, S. Z. 1965, Phys. Rev. Letters, 15, 769.
- DeShong, J. A., Hildebrand, R. H., and Meyer, P. 1964, Phys. Rev. Letters, 12, 3.

Earl, J. A. 1961, Phys. Rev. Letters, 6, 125.

Ehmert, A. 1960, Proc. Moscow Internatl. Conf. on Cosmic Rays, IV, 142.

Fan, C. Y., Gloeckler, G., and Simpson, J. A. 1965, Univ. Chicago Report,
EFINS, 65-22.

Fermi, E. 1949, Phys. Rev. 75, 1169.

Fichtel, C. E., Guss, D. E., Kniffen, D. A., and Neelakantan, K. A. 1964,
J. Geophys. Res., 69, 3293.

Freier, P. S., and Waddington, C. J. 1965, Space Science Reviews, 4, 313.

Ginzburg, V. L., and Syrovatskii, S. I. 1961, Prog. Theor. Phys. (Kyoto),
Suppl. 20, 1.

----- 1964, The Origin of Cosmic Rays, (Pergamon
Press).

Gould, R. J., and Burbidge, G. R. 1965, Preprint.

Hartman, R. C., Meyer, P., and Hildebrand, R. H. 1965, J. Geophys. Res.,
70, 2713.

Hayakawa, S., and Kitao, K. 1956, Prog. Theor. Phys., 16, 139.

Hayakawa, S., and Okuda, H. 1962, Prog. Theor. Phys., 28, 517.

Jones, F. C. 1963, J. Geophys. Res., 68, 4399.

Kraushaar, W. L., Clark, G. W., Garmire, G., Helmken, H., Higbie, P.,
and Agogino, M. 1965, Ap. J., 141, 845.

Lam, H., and Sandri, G. 1965, private communication.

L'Heureux, J., and Meyer, P. 1965, Phys. Rev. Letters, 15, 93.

McDonald, F. B., and Ludwig, G. H. 1964, Phys. Rev. Letters, 13, 783.

McDonald, F. B., and Webber, W. R. 1961, GSFC Contributions to Kyoto Conf.
on Cosmic Rays and Earth Storm, unpublished.

Meyer, P. and Vogt, R. 1961, Phys. Rev. Letters, 6, 193.

Parker, E. N. 1963, Interplanetary Dynamical Processes, (New York; Inter-
science).

Pollack, J. B., and Fazio, G. G. 1963, Phys. Rev., 131, 2684.

Schmoker, J. W., and Earl, J. A. 1965, Phys. Rev., 138, B-300.

FIGURE CAPTIONS

Figure 1: Observed cosmic-ray electron intensities in the $1 < \gamma < 10^6$ region. Most of the differential intensities shown are taken directly from the references (Cline, Ludwig and McDonald, 1964, in which the diamonds represent steady values and the crosses time variations; Schmoker and Earl, 1965; Meyer and Vogt, 1961; Earl, 1961; L'Heureux and Meyer, 1965). The integral intensities of Agrinier, et al. (1964) and Daniel and Stephens (1965) were converted to differential intensities by assuming a slope of -1.5 and an infinitely high cutoff.

Figure 2a: The low-energy electron observations plotted, for comparison, with a smooth fit to a composite of the low-energy proton and alpha data, in which the alpha intensities have been multiplied by 5. The dashed curve is an extrapolation, beyond the measurements, of the proton and alpha fit, devised by Balasubrahmanyam, Boldt and Palmeira (1965).

Figure 2b: The same data, for which the proton curve has been demodulated by the factor $\exp(K/\beta)$ with three values of K and for which the electron data have been altered with $K = 1$. Higher-energy proton fits (McDonald and Webber, 1961; Ginsburg and Syrovatskii, 1964) are shown for comparison. These curves were taken from Brunstein and Cline (1965).

Figure 3: The source strength for galactic knock-on production, Equation 5, plotted for comparison with the approximation of Equation 6. The approximation is used in all the calculations.

Figure 4: Beta-decay spectra of monoenergetic neutrons. Cosmic-ray neutrons of characteristically BeV energies are seen to produce electrons having

a spectrum in the MeV region; the source strength from galactic secondary neutrons is found, however, to have a significantly lower intensity than does that from knock-on production.

Figure 5a: Calculated differential intensities of electrons from galactic secondaries are compared with sample observations. The results are shown with and without losses due to synchrotron radiation and losses due to escape from the galaxy, using a halo value of the matter density. Also shown is the result in which the disk value of the density was used, and losses were incorporated. When no losses are used, the result is independent of the density.

Figure 5b: Upper limits to the calculated differential intensities of electrons from metagalactic secondary production. Again, synchrotron losses and red-shift escape losses are varied for comparison. In all cases, the metagalactic intensities are below the galactic intensities.

Figure 6: Calculated differential intensity of electrons resulting from galactic knock-on production and accelerated by a heliocentric field of 1.5 MV. If such a field exists, the low-energy observations can be accounted for, but the high-energy electrons remain in excess of secondary meson production.

Figure 7: Calculated differential intensities of secondary electrons which have been accelerated by a galactic Fermi process. Various values of α are incorporated for which the results are plotted in a) without, and in b) with, escape losses. The galactic halo value of the matter density is used throughout. It is seen that the observations, samples of which are shown, can be roughly fitted with a sufficiently large value of α . For certain combinations of the galactic

parameters ρ , b , λ and a , peaks in the intensity would occur at normalized energies equal to k/a .

Figure 8: Calculated differential intensities of secondaries for a fixed ratio of $\lambda/a = 1.5$. Here the singularity disappears at the high energies, but the low-energy fit becomes independent of λ and a and is seen to be inadequate to account for the observations.




Fig. 1.




Fig. 2.




Fig. 3.




Fig. 4.





Fig. 5.




Fig. 6.





Fig. 7.




Fig. 8.